

Tyron A. Alli, MD
Alexander B. Bernal, MD
John J. Cannella III, MD
Jason J. Cisler, MD
Rebecca A. Ehlers, MD
Joshua T. Evans Sr., MD
Helen Fasanya-Uptagraft, MD

Benjamin S. Hall, MD Kimberly S. Harmon, MD Jordan D. Holmes, MD Grant F. Hutchins, MD William C. Livingston, DO John C. Mitchell II, MD Matthew M. McMahon, MD Thomas R. McGinn, MI Sheeva K. Parbhu, MD Trevor J. Pearson, MD Kyle D. Rose, DO Michael E. Schafer, MD Marc A. Scheer, DO Bradley J. Schroeder, M Brian W. Ward, MD

## **Post Injectafer Instructions**

- 1. Resume diet, normal activities and medication except oral iron tablets.
- 2. Serious allergic reactions are rare however you may experience headache and nausea. Please call the office at 402-397-7057 if these symptoms are severe.
- 3. IV site: If site is tender, apply a warm washcloth for 15-20 minutes four times a day to ease the discomfort.
- 4. Follow up visit is scheduled for: FUTURE APPOINTMENTS
- 5. Four weeks after your second treatment, you will be contacted to arrange for follow-up lab work.

Additional Instructions: N/A