



Tyron A. Alli, MD
Alexander B. Bernal, MD
John J. Cannella III, MD
Jason J. Cisler, MD
Rebecca A. Ehlers, MD
Derrick D. Eichele, MD
Helen Fasanya-Uptagraft, MD
Benjamin S. Hall, MD

Kimberly S. Harmon, MD
Jordan D. Holmes, MD
Grant F. Hutchins, MD
Deepti A. Jacob, MD
William C. Livingston, DO
Thomas R. McGinn, MD
Matthew M. McMahon, MD
Sheeva K. Parbhu, MD

Trevor J. Pearson, MD
Thoetchai "Bee" Peeraphatdit, MD
Rajani Rangray, MD
Kyle D. Rose, DO
Michael E. Schafer, MD
Marc A. Scheer, DO
Brian W. Ward, MD

OMAHA | BELLEVUE | COUNCIL BLUFFS | FREMONT
402.397.7057 midwestgi.com

8901 Indian Hills Drive, Suite 200, Omaha, NE 68114

Prep Instructions

Prep Instructions

PLENVU SPLIT - 2 DAY COLON PREP INSTRUCTIONS

Procedure Details:

Procedure Date:
Procedure Check-in Time:
Procedure Time:
Procedure Location:
Scheduled with Provider:

PLEASE READ ALL INSTRUCTIONS ON THE DAY YOU RECEIVE THEM

Your provider has referred you for a colonoscopy. Bowel preparation (cleansing) is needed to perform an effective procedure. Your provider has ordered you a Plenvu split dose prep as recent studies have clearly shown a split dose prep results in higher quality examinations with increased detection of precancerous polyps. Any stool remaining in the colon can hide lesions and result in the need to repeat the colonoscopy. It is critical that you follow the prep instructions as directed below.

Contact our office immediately at 402-397-7057 if any of the following instances occur between the time you scheduled your procedure to the day of your appointment:

1. If you start a blood thinner
2. If you start any new medications
3. If you start a diabetic or weight loss medication
4. Develop COVID or upper respiratory infection
5. If you need to RESCHEDULE or CANCEL your appointment

Note: If you are of child-bearing potential, please contact your primary provider to arrange for a urine pregnancy test seven (7) days prior to your procedure. If you are certain you cannot be pregnant and prefer to refrain from prescribed testing, you may sign a pregnancy waiver the day of the procedure confirming you cannot be pregnant.

1. Your prescription for Plenvu will be electronically sent to your pharmacy 2 weeks prior to your procedure.
2. Contact your pharmacy two (2) weeks prior to your procedure to ensure your Plenvu prescription is ready for pick up.
3. Pick up your prep from your pharmacy in a timely manner or the prep may be restocked.

FIVE DAYS BEFORE YOUR COLONOSCOPY:

1. Familiarize yourself with all the colon prep instructions and contact our office at 402-397-7057 if you have any questions.
2. Purchase the following over the counter supplies:
 - Magnesium Citrate 10-ounce bottle
3. STOP medications specifically instructed to hold (only applies to patients instructed during the scheduling process).
4. Refrain from eating all types of nuts, popcorn, seeds and granola until after your procedure
5. STOP all of the following if taking:
 - Herbal supplements
 - Curcumin/Turmeric
 - Vitamins
 - Stool bulking agents (such as Metamucil or Citrucel)
 - Iron supplements
6. DO NOT STOP aspirin if you have been instructed by a physician to take daily.

TWO DAYS BEFORE YOUR COLONOSCOPY: Upon AWAKENING:

1. You may eat a low residue breakfast (see attached informational sheet)
2. After 12:00 pm: begin CLEAR LIQUID DIET ONLY
3. Clear liquids described as "a drink you can see through."
4. DO NOT consume liquids that are red, purple or blue in color.
5. Drink clear liquids every hour throughout the day from the following list:
 - Clear fruit juices (white grape or apple juice)

- Water, tea, or coffee (without cream)
- Kool-Aid or PowerAde
- Clear soup, broth, or bouillon
- Popsicles
- Hard candies
- Soda: regular or diet (7-Up, Sprite, Pepsi, Coke, Ginger Ale, Orange).
- Jell-O

Note: If you are a diabetic patient, consume non-sugar-free clear liquids for calorie support in blood sugar management
6. NO SOLID FOOD, ALCOHOL, or RECREATIONAL DRUGS.

7. Drink 10-ounces Magnesium Citrate between 6:00 pm and 7:00 pm.

THE DAY BEFORE YOUR COLONOSCOPY: Upon AWAKENING:

1. Continue CLEAR LIQUID DIET (see list above).
2. NO SOLID FOOD, ALCOHOL, or RECREATIONAL DRUGS.
3. Stay hydrated: drink at least 8-ounces of clear liquids every hour

At 5:00 pm -take your first dose of Plenvu prep as follows:

- Using the supplied container, mix contents (dose 1 - Mango flavor) with at least 16-ounces of water
- Mix the contents by shaking or using a spoon until completely dissolved (2-3 minutes).
- Drink ALL the Plenvu mixture in the container over 30 minutes.
- Refill container with at least 16-ounces of any clear liquid and drink over the next 30 minutes.

Note: continue to drink clear liquids until you go to bed.

THE DAY OF YOUR COLONOSCOPY:

1. NO SOLID FOOD, ALCOHOL, OR RECREATIONAL DRUG USE.
2. NO CHEWING/SMOKELESS TOBACCO PRODUCTS.

Start second part of prep: At [Six hours prior] six hours prior to arrival & check-in, begin drinking 2nd half of Plenvu prep.

- Repeat steps above using supplied container.
- Mix contents of Dose two (2) (pouch A & B - fruit punch flavor) with 16-ounces of water.
- Mix the contents by shaking or using a spoon until completely dissolved (2-3 minutes).
- Drink ALL the Plenvu mixture in the container over 30 minutes.
- Refill container with at least 16-ounces of any clear liquid and drink over the next 30 minutes.
- You must be completely finished with your prep 4 hours prior to arrival at the facility.
- TAKE NOTHING ELSE BY MOUTH (NO EATING, NO DRINKING) AFTER YOU HAVE FINISHED THE PREP.
- Your bowel movements should be clear-yellow liquid (urine or lemonade looking).
- If your bowel movements are still brown and not clear, please call our office at 402-397-7057 (24hr service).
- Take your morning medications before 6:00 am with a few sips of water.

Note: If you are a DIABETIC PATIENT, please refer to "Special Instruction" sheet sent.

RIDE HOME:

You must have a responsible driver who has a valid driver's license to take you home. It is preferred that you have someone drive you to the scheduled location, wait while you have your procedure, and then accept responsibility for your dismissal upon leaving the facility. In the event that your driver is not present at check-in you will be expected to validate your responsible driver. If check-in staff are unable to validate your responsible driver, your colonoscopy will be cancelled.

IMPORTANT: Public transportation (including rides shares such as Uber or Lyft) is NOT an acceptable form of transportation following your procedure unless accompanied by a responsible adult. If you are scheduled at a hospital, please know individual hospital policies may require you to have a responsible adult stay with you for 24 hours post procedure.

FOLLOWING YOUR COLONOSCOPY:

1. Drink 8-ounces of liquid six (6) times before retiring for the night.
2. Do NOT drive, operate machinery, return to work or make important decisions for the remainder of the day.
3. You may resume normal activities the next day unless your provider states otherwise.

WHAT TO BRING TO YOUR COLONOSCOPY:

1. If you were sent forms through the patient portal, please complete them prior to your visit,
2. Be prepared to provide first and last name of all your providers you want to receive a copy of your procedure report.
3. Responsible adult to drive you home (NO EXCEPTIONS). If no responsible adult driver, your procedure will be cancelled.
4. Insurance card(s)
5. Photo ID

WHAT TO WEAR TO YOUR COLONOSCOPY:

1. Wear comfortable, loose-fitting clothing.
2. Wear flat or tennis shoes.
3. Please leave jewelry and valuables at home.

FREQUENTLY ASKED QUESTIONS:

Q. Why do I have to drink my prep at two different times?

A. Recent studies have clearly shown split dose preps result in higher quality examinations with increased detection of precancerous polyps.

Q. What if I feel like I am going to vomit if I drink another glass of prep or what if I start to vomit while drinking the prep?

A. Stop drinking the prep for 30-45 minutes until symptoms subside, then resume prep.

Q. What if I drink all of the prep and I do not have a bowel movement?

A. If no stool by 9:00 pm, you should use a fleets enema. This can be purchased over the counter at any pharmacy. If after you administer the enema you still have no results, call our office at 402-397-7057 (24hr line).

Q. If I have clear colored stool return after consuming the 1st half of the prep, do I have to drink the 2nd half?

A. Yes, it is important that you drink the entire prep. Digestion continues throughout the night requiring the 2nd half of the prep to be drank to ensure best results.

Q. What should my bowel movements look like after I drink all of the prep solution?

A. Your bowel movements should be clear yellow liquid. They will look like urine or lemonade. If your bowel movements are still brown and haven't been clear, please call our office at 402-397-7057 (24hr line).

Q. What if I have rectal discomfort?

A. You may apply petroleum-based product or diaper rash ointment to the rectal area if you experience discomfort from frequent stools.

Q. Does my driver have to stay with me during the procedure?

A. If your driver does not wish to remain in the lobby, a contact number can be given to the nursing staff. These arrangements must be made during the check-in process, or your procedure will be cancelled. Typically, the driver can return two hours after they drop you off or they can be called prior to your dismissal time. Public transportation can only be used if you are accompanied by a responsible adult.

Q. Can I eat after my procedure?

A. Usually you may resume your normal diet unless otherwise instructed. Q. Will I be asleep for my procedure? A. Yes, sedation will be provided to keep you comfortable throughout your procedure(s).

Q: Will my insurance pay for my colonoscopy?

A: If you are scheduled for a colonoscopy: Please understand all colonoscopies are not screening/preventative and may be deemed medical based on your medical history. Deductibles, copays and co-insurance may apply. The Facility submits procedural documentation and charges according to the Medicare and Medicaid services and American Medical Association guidelines and is not responsible for determining how benefits are paid by your insurance plan. Please make sure your insurance company allows your procedure at our Facility. If you have questions, please call us at (402) 504-3846 for a more detailed explanation.

To bill for a screening colonoscopy the patient must be without gastrointestinal symptoms, 45 years of age or older, have no personal history of gastrointestinal disease, colon polyps and/or colon cancer. In most cases patients are limited to one screening colonoscopy per ten-year increment.